



What does the research tell us about the potential long-term health effects from repeated head impacts and concussion?

We are aware that people are hearing mixed messages about the potential long-term effects from concussion and repeated head impacts. To address this, we have worked with experts across New Zealand to develop this summary for you.

What do we mean by concussion and repeated head impacts?

Concussion is a form of mild traumatic brain injury. The person may experience dizziness or confusion immediately after the injury. Symptoms can include headaches, sensitivity to light, feeling extremely tired, or difficulty remembering things.

Repeated head impacts are where a person experiences multiple hits to the head or body that can cause the brain to move within the hard skull e.g., from a hard tackle or an assault. The person may not have any immediate symptoms, but the effects may add up over time.

What are the potential long-term impacts?

1. Development of Brain Disorders – There is consistent evidence showing a link between concussion and increased risk of developing brain disorders such as dementia and stroke.¹⁻⁵ However, the size of this risk (e.g., how many people will be affected and how many concussions is too many) remains unclear. Concussion is one of 14 known risk factors for dementia.⁶

2. Cognitive Functioning Difficulties – There is consistent evidence showing a link between repeated concussions (e.g., lots and/or in quick succession) and difficulties in a person's ability to think later in life. The cognitive functions that can be affected include: difficulties making decisions quickly, being able to multi-task, remembering things, processing new information and reduced self-control.⁷⁻⁹ We still do not know how big the risk is or how many concussions it takes to cause long-term cognitive issues. For the majority of people, experiencing a few concussions over your lifetime does not impact your ability to think in later life.¹⁰

3. Chronic Traumatic Encephalopathy (CTE) – CTE is a term that describes degenerative changes in the brain's structure. CTE is identified by clusters of a protein called tau that form in specific areas of the brain. These tau clusters can only be seen following a medical exam of the brain after death (autopsy).^{11,12} Research indicates there is an association between repeated head impacts sustained over many years in contact sports and violence, and increased risk of developing CTE.¹³⁻¹⁶ We still do not know how the tau clusters in the brain relate to symptoms experienced during life or how much of the tau protein in the brain is problematic. The symptoms people may experience in later life have not been linked exclusively to CTE changes seen in the brain.

Many of the symptoms that people experience following concussion and repeated head impacts can be treated to reduce the risk of long-term problems.¹⁷⁻²⁰



References

1. Mackay DF, Russell ER, Stewart K, MacLean JA, Pell JP, Stewart W. Neurodegenerative Disease Mortality among Former Professional Soccer Players. *N Engl J Med* 2019; 381(19): 1801-8.
2. Barnes DE, Byers AL, Gardner RC, Seal KH, Boscardin WJ, Yaffe K. Association of Mild Traumatic Brain Injury With and Without Loss of Consciousness With Dementia in US Military Veterans. *JAMA Neurol* 2018; 75(9): 1055-61.
3. Fann JR, Ribe AR, Pedersen HS, et al. Long-term risk of dementia among people with traumatic brain injury in Denmark: a population-based observational cohort study. *Lancet Psychiatr* 2018; 5(5): 424-31.
4. Turner GM, McMullan C, Aiyegbusi OL, et al. Stroke risk following traumatic brain injury: Systematic review and meta-analysis. *Int J Stroke* 2021; 16(4): 370-84.
5. Anns F, Quarrie KL, Milne BJ, et al. Neurodegenerative Diseases in Male Former First-Class New Zealand Rugby Players. *Sports Med* 2025.
6. Livingston G, Huntley J, Liu KY, et al. Dementia prevention, intervention, and care: 2024 report of the Lancet standing Commission. *The Lancet* 2024; 404(10452): 572-628.
7. Gallo V, Motley K, Kemp SPT, et al. Concussion and long-term cognitive impairment among professional or elite sportpersons: a systematic review. *J Neurol, Neurosurg Psychiatr* 2020; 91(5): 455-68.
8. Manley G, Gardner AJ, Schneider KJ, et al. A systematic review of potential long-term effects of sport-related concussion. *Br J Sports Med* 2017; 51(12): 969-77.
9. Hume P, Theadom A, Lewis G, et al. A Comparison of Cognitive Function in Former Rugby Union Players Compared with Former Non-Contact-Sport Players and the Impact of Concussion History. *Sports Med* 2017; 47(6): 1209-20.
10. Theadom A, Barker-Collo S, Parag V, et al. Mild Traumatic Brain Injury Does Not Significantly Affect Midlife Cognitive Functioning Within the General Population: Findings From a Prospective Longitudinal Birth Cohort Study. *J Head Trauma Rehabil* 2024; 39(2): E70-e82.
11. Bieniek KF, Cairns NJ, Crary JF, et al. The Second NINDS/NIBIB Consensus Meeting to Define Neuropathological Criteria for the Diagnosis of Chronic Traumatic Encephalopathy. *J Neuropathol Exp Neurol* 2021; 80(3): 210-9.
12. McKee AC, Stein TD, Huber BR, et al. Chronic traumatic encephalopathy (CTE): criteria for neuropathological diagnosis and relationship to repetitive head impacts. *Acta Neuropathol* 2023; 145(4): 371-94.
13. Daneshvar DH, Nair ES, Baucom ZH, et al. Leveraging football accelerometer data to quantify associations between repetitive head impacts and chronic traumatic encephalopathy in males. *Nature Comm* 2023; 14(1): 3470.
14. Bieniek KF, Blessing MM, Heckman MG, et al. Association between contact sports participation and chronic traumatic encephalopathy: a retrospective cohort study. *Brain Pathology* 2020; 30(1): 63-74.
15. Mez J, Daneshvar DH, Abdolmohammadi B, et al. Duration of American Football Play and Chronic Traumatic Encephalopathy. *Ann Neurol* 2020; 87(1): 116-31.
16. Nowinski CJ, Bureau SC, Buckland ME, et al. Applying the Bradford Hill Criteria for Causation to Repetitive Head Impacts and Chronic Traumatic Encephalopathy. *Front Neurol* 2022; 13: 938163.
17. Moore BM, Stark RK, D'Angelo EC. Multidisciplinary care for patients with persistent symptoms following concussion: a systematic review. *Disabil Rehabil* 2024; 46(9): 1760-75.
18. Möller MC, Lexell J, Ramsay KW. Effectiveness of specialized rehabilitation after mild traumatic brain injury: A systematic review and meta-analysis. *J Rehabil Med* 2021; 53(2): jrm00149.
19. Heslot C, Azouvi P, Perdrieau V, Granger A, Lefèvre-Dognin C, Cogné M. A Systematic Review of Treatments of Post-Concussion Symptoms. *J Clin Med* 2022; 11(20).
20. Silverberg ND, Lee K, Mikolic A, et al. Active collaborative on traumatic brain injury care: adapted clinical practice guideline. *J Neurotrauma*, 2025; doi: 10.1177/08977151378894.

With thanks to; Professor Alice Theadom, Professor Patria Hume, Sir Richard Faull, Dr Helen Murray, Dr Josh McGeown, Professor Lynette Tippett, Professor Maurice Curits, Dr Rosamund Hill, Dr Josh Faulkner, Professor Melanie Bussey, Associate Professor Mangor Pedersen, Professor Daniel Shepherd, Professor Richard Siegert, Katherine Forch, Dr Dion Henare, Dr Sharon Olsen.

NOTE: We acknowledge that there are research studies that show different results to those described in this statement. Differences often occur due to different populations being studied and differences in research methods used. We believe on the weight of the current evidence that these statements reflect the overall message that can be determined from the overall evidence base at this point in time. We will continue to update this summary as more information becomes available.

