



## Brain Injury Screening Tool (BIST-3)

### A guide for traumatic brain injury assessment

The Brain Injury Screening Tool (BIST-3) is a brief tool for use on initial presentation for suspected mild traumatic brain injury/ concussion.

Its purpose is to help guide the clinical assessment by operationalising current international best practice guidelines.<sup>1</sup>

The BIST was developed for health professionals working across primary and secondary health care.

The BIST facilitates clinical decision making through identification of people who are at risk of longer-term difficulties who may benefit from early specialist treatment.

This tool should be used in addition to clinical judgment and other assessments such as the Vestibular/Oculomotor Motor Screening (VOMS), King-Devick or the Romberg's test. Additional questioning to add to the clinical picture is encouraged.

The first 10 questions can be administered in any order to assist flow of the consultation. The symptom scale should be completed as presented.

The symptom scale and impact item can be repeated at follow up to monitor recovery over time.

Date of Injury (dd/mm/yy):

Time of Injury (HH:MM 24hr):

Date of Consultation (dd/mm/yy):

Age:

Gender:

Ethnicity:

1. Please tell me what happened:

2. Are there high risk indicators such as suspicion of skull fracture, focal neurological deficit, high speed, focal blunt trauma or fall from height? (e.g. >5 stairs)

Yes No

\*If high risk factors are present consider referral to the ED or further evaluation in hospital

3. Did the incident occur in traumatic circumstances which could result in emotional or psychological reactions? (e.g., assault, domestic violence, fatalities in a car accident)

Yes No

4. Did anyone with you at the time of the injury say anything else about what happened?

Yes No

5. Were you sick or did you vomit?

Yes No a. If yes, how many times

6. Were you knocked out (or did you lose consciousness)?

Yes No Unknown a. If yes, how long hrs mins

7. Are you feeling better, worse or about the same since the injury?

Better Worse About the same

8. Have you had a concussion or brain injury before?

Yes No If yes, (i) how many times?

(ii) When was the last injury?(dd/mm/yy)

(iii) Did your last injury take more than 3 months to recover from? Yes No

9. Have you ever experienced any difficulties with your mental health?

Yes No

10. Do you have a history of migraine? (severe headache with vomiting or extreme sensitivity to light and sound)

Yes No


11. Please ask the patient the following question.

Compared with before the accident, please rate how much you experience the following symptoms:

												
		Not at all	Mild (a little)			Moderate (quite bad)				Severe (very bad)		
		0	1	2	3	4	5	6	7	8	9	10
Physical	Headache (my head hurts)											
	My neck hurts											
	I don't like bright lights											
	I don't like loud noises											
Total physical score (out of 40)												
Vestibular-ocular	I feel dizzy or like I could be sick											
	If I close my eyes, I feel like I am at sea											
	I have trouble with my eyesight (vision)											
	I feel clumsy (bumping into things or dropping things more than usual)											
Total vestibular score (out of 40)												
Cognitive	It takes me longer to think											
	I forget things											
	I get confused easily											
	I have trouble concentrating											
Total cognitive score (out of 40)												
If more than 24 hours post-injury, please also rate these physical symptoms												
	I get angry or irritated easily											
	I just don't feel right											
	I feel tired during the day											
	I need to sleep a lot more or find it hard to sleep at night											

12. Injuries to the brain can affect how a person feels, behaves, thinks and how able they are to do everyday tasks.

On a scale of 0 to 10, how much do you feel your injury is impacting on you? Where 0 means that the injury has not had any impact on you and 10 means you feel that injury impacts on everything you do.



**Please check if the person needs to go to the Emergency Department or further evaluation in hospital (e.g., suspected skull fracture, severe headache, use of blood anti-coagulants) before considering whether a referral to rehabilitation service is required.**

Total symptom severity score  
less than 24 hours (out of 120)

Total symptom severity score  
greater than 24 hours (out of 160)

Overall score

**What is the dominant symptom cluster?**  
(High proportion or most severe symptoms  
reported; e.g., physical, vestibular or cognitive?)

Total physical score: /40  
Total vestibular score: /40  
Total cognitive score: /40

If the dominant symptom cluster is Vestibular or if a vestibular item is rated >8 at 7-10 days post-injury, consider referral to a physiotherapist or Concussion Service.

**Community referral recommendation**

(please ensure all questions have been answered to obtain an accurate recommendation)

**Acknowledgements**

This tool has been developed by the BIST Development Group: Alice Theadom, Natalie Hardaker, Penelope Day, Kris Fernando, Katherine Forch, Kevin Henshall, Doug King, Michelle Wilkinson, Renata Bastos-Gottgroy, Sam Jewell, Stephen Kara, Jason Chua, Patria Hume.

**Reference**

Silverberg ND, et al on behalf of the American Congress of Rehabilitation Medicine Brain Injury Interdisciplinary Special Interest Group Mild TBI Task Force.

Management of Concussion and Mild Traumatic Brain Injury: A Synthesis of Practice Guidelines. Archives of Physical Medicine and Rehabilitation, 2020, 101; 382-393

© 2022. BIST Development Group. The Brain Injury Screening Tool (BIST) is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.

 CC BY-ND: This license allows reusers to copy and distribute the material in any medium or format in unadapted form only, and only so long as attribution is given to the creator. The license allows for commercial use.



**AUT TRAUMATIC BRAIN  
INJURY NETWORK**